

CORRECTION

Open Access



Correction to: To develop a regional ICU mortality prediction model during the first 24 h of ICU admission utilizing MODS and NEMS with six other independent variables from the critical care information system (CCIS) Ontario, Canada

Raymond Kao^{1,2,3*}, Fran Priestap² and Allan Donner²

Correction to: J Intensive Care

<https://doi.org/10.1186/s40560-016-0143-6>

In the original publication of this article [1], there were several transcription errors in the logistic regression equation model as below:

a. Equation published:

Log [Mortality (at 24 h ICU admission)] = $-5.18 + 0.80[\text{age}(40-79)] + 1.71[\text{age}(> 80)] + 0.60[\text{Sex}(\text{male} = 0 \text{ and female} = 1)] + 0.98[\text{Other source admission}] + 0.00[\text{Operating room/post-anesthesia care}] + 1.00[\text{ER admission}] + 1.12[\text{Hospital-outside or within LHIN}] + 1.60[\text{Ward admission}] + 0.00[\text{Cardiovascular/Cardiac/Vascular}] + 0.00[-0.81[\text{Other diagnosis}] - 0.80[\text{Gastrointestinal}] - 0.56[\text{Respiratory}] - 0.32[\text{Trauma}] + 0.002[\text{Neurological}] - 0.30[\text{ICU re-admission}] - 0.21[\text{CCI}(1-3)] + 0.05[\text{CCI}(> 3)] + 0.0[\text{NEMS}(0-22)] + 0.39[\text{NEMS}(23-29)] + 1.02[\text{NEMS}(\geq 30)] + 1.18[\text{MODS}(1-4)] + 1.91[\text{MODS}(5-8)] + 2.90[\text{MODS}(9-120)] + 3.56[\text{MODS}(\geq 130)].$

b. Corrected equation:

Logit [Mortality (at 24 h ICU admission)] = $-5.41 + 0.00[\text{age}(0-39)] + 0.80[\text{age}(40-79)] + 1.71[\text{age}(> 80)] - 0.06[\text{Sex}(\text{male} = 0 \text{ and female} = 1)] + 0.98[\text{Other source admission}] + 0.00[\text{Operating room/post anesthesia care}] + 1.00[\text{ER admission}] + 1.12[\text{Hospital-outside or within LHIN}] + 1.60[\text{Ward admission}] + 0.00[\text{Cardiovascular/Cardiac/Vascular}] - 0.81[\text{Other diagnosis}] - 0.80[\text{Gastrointestinal}] - 0.56[\text{Respiratory}] - 0.32[\text{Trauma}] + 0.002[\text{Neurological}] + 0.30[\text{ICU re-admission}] - 0.21[\text{CCI}(1-3)] + 0.05[\text{CCI}(> 3)] + 0.0[\text{NEMS}(0-22)] + 0.39[\text{NEMS}(23-29)] + 1.02[\text{NEMS}(\geq 30)] + 1.18[\text{MODS}(1-4)] + 1.91[\text{MODS}(5-8)] + 2.90[\text{MODS}(9-12)] + 3.56[\text{MODS}(\geq 13)].$

The authors sincerely apologize for the inconvenience caused to the readers.

Author details

¹Department of National Defense, Royal Canadian Medical Services, 1745 Alta Vista Drive, Ottawa, Ontario K1A 0K6, Canada. ²London Health Sciences Center, Divisions of Critical Care and Roberts Research Institute, Western University, 800 Commissioner's Rd E, London, Ontario N6A 5W9, Canada. ³Harvard School of Public Health, Harvard University, 677 Huntington Ave, Boston, MA 02115, USA.

Published online: 14 January 2020

Reference

1. Kao, et al. *J Intensive Care*. 2016;4:16.

The original article can be found online at <https://doi.org/10.1186/s40560-016-0143-6>

* Correspondence: rkao3@uwo.ca

¹Department of National Defense, Royal Canadian Medical Services, 1745 Alta Vista Drive, Ottawa, Ontario K1A 0K6, Canada

²London Health Sciences Center, Divisions of Critical Care and Roberts Research Institute, Western University, 800 Commissioner's Rd E, London, Ontario N6A 5W9, Canada

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.