

CORRECTION

Open Access



Correction to: The development and validation of a “5A” severity scale for predicting in-hospital mortality after accidental hypothermia from J-point registry data

Yohei Okada^{1,2*}, Tasuku Matsuyama³, Sachiko Morita⁴, Naoki Ehara⁵, Nobuhiro Miyamae⁶, Takaaki Jo⁷, Yasuyuki Sumida⁸, Nobunaga Okada^{3,9}, Makoto Watanabe³, Masahiro Nozawa¹⁰, Ayumu Tsuruoka¹¹, Yoshihiro Fujimoto¹², Yoshiki Okumura¹³, Tetsuhisa Kitamura¹⁴, Shungo Yamamoto¹⁵, Ryoji Iiduka² and Kaoru Koike¹

Correction to: J Intensive Care
<https://doi.org/10.1186/s40560-019-0384-2>


In the original publication of this article [1], the figure legends of Fig. 2 and Fig. 3 are wrong and should be changed as below:

5A scoring model			Severity	Sum Risk score	Predicted mortality*
A ge	Covariates	Score			
		60-69	Low risk	0-3	7.1 % [5.8-8.4]
		70-79	Mild	4	20.5 % [18.7-22.3]
ADL		≥ 80	Moderate	5	35.4 % [33.3-37.6]
	Disturbance	1	Severe	6-9	67.4 % [65.1-69.6]
A rrest		Arrest or SBP ≤ 60 mmHg	*Predicted mortality: Mean [95%CI] in the validation cohort		
A cidemia		pH: 7.2-7.35			
		<7.2			
A lbumin		≤ 3 (mg/dl)			
Sum score					/9

Fig. 2 5A scoring model

* Correspondence: yokada-kyf@umin.ac.jp
¹Department of Primary Care and Emergency Medicine, Graduate School of Medicine, Kyoto University, 606-8501, Yoshidakonoe-cho, Sakyo, Kyoto, Japan
²Department of Emergency and Critical Care Medicine, Japanese Red Cross Society, Kyoto Daini Hospital, Kyoto, Japan
 Full list of author information is available at the end of the article





Author details

¹Department of Primary Care and Emergency Medicine, Graduate School of Medicine, Kyoto University, 606-8501, Yoshidakonoe-cho, Sakyo, Kyoto, Japan. ²Department of Emergency and Critical Care Medicine, Japanese Red Cross Society, Kyoto Daini Hospital, Kyoto, Japan. ³Department of Emergency Medicine, Kyoto Prefectural University of Medicine, Kyoto, Japan. ⁴Senri Critical Care Medical Center, Saiseikai Senri Hospital, Suita, Japan. ⁵Department of Emergency, Japanese Red Cross Society, Kyoto Daiichi Red Cross Hospital, Kyoto, Japan. ⁶Department of Emergency Medicine, Rakuwa-kai Otowa Hospital, Kyoto, Japan. ⁷Department of Emergency Medicine, Uji-Tokushukai Medical Center, Uji, Japan. ⁸Department of Emergency Medicine, North Medical Center, Kyoto Prefectural University of Medicine, Kyoto, Japan. ⁹Department of Emergency and Critical Care Medicine, National Hospital Organization, Kyoto Medical Center, Kyoto, Japan. ¹⁰Department of Emergency and Critical Care Medicine, Saiseikai Shiga Hospital, Ritto, Japan. ¹¹Department of Emergency and Critical Care Medicine, Kyoto Min-Iren Chuo Hospital, Kyoto, Japan. ¹²Department of Emergency Medicine, Yodogawa Christian Hospital, Osaka, Japan. ¹³Department of Emergency Medicine, Fukuchiyama City Hospital, Fukuchiyama, Japan. ¹⁴Division of Environmental Medicine and Population Sciences, Department of Social and Environmental Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan. ¹⁵Department of Healthcare Epidemiology, School of Public Health in the Graduate School of Medicine, Kyoto University, Kyoto, Japan.

Published online: 11 June 2019

Reference

- Okada Y, et al. The development and validation of a "5A" severity scale for predicting in-hospital mortality after accidental hypothermia from J-point registry data. *J Intensive Care*. 2019;7:27 <https://doi.org/10.1186/s40560-019-0384-2>.